Affix patient label here

Patient's Study Participant Number:



Sampling Instructions

- The research team will inform you of the planned sample collection on the day
- Avoid suctioning or using saline washes: 1 hour before if patient has high secretion load or 2 hours if patient has a low secretion load
- Sample collection should be at least 1 hour clear from nebuliser therapy or saline washes
- If suctioning is clinically indicated please continue and inform the research team who will delay collection

Collection Day	Date	Planned collection time	Avoid routine suctioning for:	Sample collection	Planned reattempt time	Sample Collection	Send to
Monday			1 hour prior 2 hours prior	Subglottic Time: Tracheal Time:		Successful Unable Successful Unable Ime:	Direct
Tuesday			1 hour prior 2 hours prior	Subglottic Time: Tracheal Time:		Successful Unable Successful Unable Ime:	Direct
Wednesday			1 hour prior 2 hours prior	Subglottic Time: Tracheal Time:		Successful Unable Successful Unable Ime:	Direct
Thursday			1 hour prior 2 hours prior	Subglottic Time: Tracheal Time:		Successful Unable Successful Unable Ime: Time: Time:	Direct

Collection Day	Date	Planned collection time	Avoid routine suctioning for:	Sample collection	Planned reattempt time	Sample Collection	Send to
Friday			1 hour prior 2 hours prior	Subglottic Time: Tracheal Time:		Successful Unable Successful Unable Ime:	Store in fridge
Saturday			1 hour prior 2 hours prior	Subglottic Time: Tracheal Time:		Successful Unable Successful Unable Ime:	Store in fridge
Sunday			1 hour prior 2 hours prior	Subglottic Time: Tracheal Time:		Successful Unable Successful Unable Ime: Successful Unable Ime:	Store in fridge